

Research papers

Coordination
Marie-Noëlle
WOILLEZ

Authors
Lykke E. ANDERSEN
Fabiana ARGANDOÑA
Carla OLMOS
Diego CALDERON
Sebastián MIRANDA
Alvaro MUÑOZ
Sergio CHOQUE

Assessing CO2 Emissions from Deforestation and Fires in Bolivia Between 2010 and 2023

Introduction	4
1. Data and Methodology	6
1.1. General workflow	6
1.2. Deforestation emissions accounting method	7
1.2.1. Bookkeeping models	7
1.2.2. Carbon Pools	8
1.2.4. Assumptions for deforestation estimations	13
1.3. Assessment of burned area and fire emissions data	17
2. Results	21
2.1. Carbon emissions from land-use change – Deforestation	21
2.1.1. Gross emissions	21
2.1.2. Gross removals	23
2.1.3. Net fluxes	24
2.2. Burned area and fire emissions	28
2.2.1. Burned area	28
2.2.2. Fire emissions estimates 2010 – 2022	29
2.2.3. Fire emissions from forest degradation	31
2.3. National results	34
2.4. Subnational level results	35
2.4.1. Departmental level	35
2.4.2. Municipal level	37
2.5. Sensitivity analysis	39
3. Discussion	41
4. Conclusions	44
Bibliography	47

Agence française de développement

Papiers de recherche

Les *Papiers de Recherche de l'AFD* ont pour but de diffuser rapidement les résultats de travaux en cours. Ils s'adressent principalement aux chercheurs, aux étudiants et au monde académique. Ils couvrent l'ensemble des sujets de travail de l'AFD : analyse économique, théorie économique, analyse des politiques publiques, sciences de l'ingénieur, sociologie, géographie et anthropologie. Une publication dans les *Papiers de Recherche de l'AFD* n'en exclut aucune autre.

Les opinions exprimées dans ce papier sont celles de son (ses) auteur(s) et ne reflètent pas nécessairement celles de l'AFD. Ce document est publié sous l'entière responsabilité de son (ses) auteur(s) ou des institutions partenaires.

Research Papers

AFD Research Papers are intended to rapidly disseminate findings of ongoing work and mainly target researchers, students and the wider academic community. They cover the full range of AFD work, including: economic analysis, economic theory, policy analysis, engineering sciences, sociology, geography and anthropology. *AFD Research Papers* and other publications are not mutually exclusive.

The opinions expressed in this paper are those of the author(s) and do not necessarily reflect the position of AFD. It is therefore published under the sole responsibility of its author(s) or its partner institutions.

Assessing CO₂ Emissions from Deforestation and Fires in Bolivia Between 2010 and 2023

Authors

Lykke E. ANDERSEN

SDSN Bolivia

Fabiana ARGANDOÑA

SDSN Bolivia

Carla OLMOS

SDSN Bolivia

Diego CALDERON

SDSN Bolivia

Sebastián MIRANDA

SDSN Bolivia

Alvaro MUÑOZ

SDSN Bolivia

Sergio CHOQUE

SDSN Bolivia

Coordination

Marie-Noëlle WOILLET (AFD)

Abstract

This study estimates net annual CO₂ emissions from deforestation and fires in Bolivia from 2010 to 2023, considering both gross emissions and absorptions resulting from land clearing, land use change, fires, and forest regeneration. Using high-resolution annual land cover maps from MapBiomass Bolivia (1985–2023), a global biomass density map and a carbon bookkeeping model, we track carbon changes at a 100×100m resolution. To estimate emissions from forest degradation due to fires, we used the Global Fires Emissions Database and subtracted emissions from deforestation within burned areas to prevent double counting. Our results indicate that net CO₂ emissions from deforestation and gross emissions from forest degradation due to fires are annually about 70 million tCO₂ and 126 million tCO₂ respectively, corresponding to per capita emissions of 6.6 and 12.3 tCO₂/pers./year.

Keywords

Deforestation; fires; carbon emissions; Bolivia.

Acknowledgements

This research has benefited from the financial support of AFD.

Original version

English

Accepted

November 2025

Résumé

Cette étude estime les émissions nettes annuelles de CO₂ issues de la déforestation et des incendies en Bolivie entre 2010 et 2023, en prenant en compte à la fois les émissions brutes et les absorptions résultant du défrichement, des changements d'usage des terres, des incendies et de la régénération forestière. En utilisant des cartes annuelles d'occupation des sols à haute résolution issues de MapBiomass Bolivia (1985–2023), une carte mondiale de densité de biomasse et un modèle de comptabilité carbone, nous suivons les changements de stocks de carbone à une résolution de 100×100m. Pour estimer les émissions liées à la dégradation des forêts par les incendies, nous avons utilisé la base de données Global Fire Emissions Database et soustrait les émissions liées à la déforestation survenues dans les zones brûlées afin d'éviter les doubles comptages. Nos résultats montrent que les émissions nettes de CO₂ dues à la déforestation et les émissions brutes résultant de la dégradation des forêts par les incendies s'élèvent annuellement à environ 70 millions et 126 millions de tonnes de CO₂ respectivement, ce qui correspond à des émissions par personne de 6.6 et 12.3 tCO₂/pers./an.

Mots-clés

Déforestation; feux; émissions de carbone ; Bolivie.

Remerciements

Ce travail de recherche a bénéficié du soutien financier de l'AFD.

Version originale

Anglais

Acceptée

Novembre 2025

Introduction

Bolivia has approximately 55 million hectares of forests, covering nearly half of its national territory. These ecosystems represent not only a strategic reserve of environmental services (Andersen et al., 2025) but also a critical global carbon sink. However, in recent years, the country has experienced rising deforestation and wildfires, leading to significant carbon emissions.

According to official reports, land-use change is the country's largest source of CO₂ emissions, accounting for over 50% of total emissions (APMT, 2020). Consequently, Bolivia ranks among the top 20 countries for land-use change emissions for 2010–2022 (Friedlingstein et al., 2023). Further, the Global Carbon Atlas in 2023 placed Bolivia as the 10th highest emitter from land-use change, 3rd in per capita emissions for this sector, and 81st in CO₂ emissions from fossil fuels (Global Carbon Atlas, 2023).

The Agriculture, Forestry, and Other Land Use (AFOLU) sector remains one of the few that includes greenhouse gas (GHG) absorption processes. However, some national inventories — particularly in data-limited countries — simplify these dynamics by assuming static scenarios or ignoring ecosystem recovery post-deforestation.

These recovering areas not only recapture part of previously released carbon but also highlight the dynamism of Bolivia's forest landscapes. On the other hand, estimating fire emissions remains fraught with technical and methodological challenges (Viglione, 2023). Based on the available data, this study includes some insights and a preliminary gross estimation of emissions resulting from vegetation degradation by fires.

In order to provide new estimates of CO₂ emissions from both deforestation and fires in Bolivia, we employ a dynamic modeling approach based on a pixel-level carbon bookkeeping model. This method tracks carbon flows by forest type, land-use change category, year accounting deforestation and forest regrowth, while incorporating changes in carbon pools (aboveground biomass, belowground biomass, and soil organic carbon).

Our study builds upon the foundational work of Andersen et al. (2016), who developed a carbon bookkeeping model to estimate net CO₂ emissions from deforestation in Bolivia during 1990–2000 and 2000–2010. Their methodology, which accounted for deforestation at a 10×10 km resolution, provided critical insights into spatial and temporal variations in emissions across Bolivia's diverse forest types.

Their pioneering work revealed rising emissions and consistently high per capita CO₂ outputs but faced limitations in

the spatial and temporal granularity of the data and excluded fire-induced degradation. Our research advances this framework by incorporating high-resolution land-cover transitions from MapBiomas with 30×30 m resolution from 2010 to 2023 and integrating fire emission data from GFED, enabling pixel-level tracking and explicit quantification of degradation

1. Data and Methodology


1.1. General workflow

This study primarily focuses on estimating deforestation CO₂ emissions based on the principles of bookkeeping models. In addition, we conduct an exploratory analysis of fire-related emissions, offering insights that contribute to the broader discussion on this topic. The following steps outline the general workflow.

- **Step 1. Establish biomass baseline by forest type.** The first step is to establish a baseline for forest biomass by forest type to estimate emissions and removals within these diverse ecosystems.
- **Step 2. Estimate carbon emissions from land-use change and forest loss.** Subsequently, an assessment of land cover and land cover transition data will be conducted to establish a timeline for annual emission estimations, primarily focusing on deforestation and conversion to other land uses. Following the principles of bookkeeping models, land use and land-use change will be central to the emissions analysis. This approach will provide an overview of forest loss, gain, and dynamics.
- **Step 3. Assess fire data of burned area and fire emissions.** The next step in analysing and understanding land-use dynamics will be the assessment of fire data and fire emissions data.
- **Step 4. Aggregate results.** Finally, aggregated results will be estimated for the country. Given the data treatment, analysis, and research decisions made in the preceding steps, gross emissions are estimated using the principles of bookkeeping models. This involves implementing response curves for different carbon pools to estimate gross and net emissions and removals. Deforestation analysis is performed at a 100-meter grid scale, while fire analysis uses a 500-meter grid scale. For reporting purposes, aggregated data for Bolivia, as well as aggregated results by department and municipality, are calculated.
- **Step 5. Sensitivity analysis.** By varying certain assumptions, we explore alternative outcomes.

Figure 1 presents the relation between data sources and processes.

Figure 1. General flowchart, data and steps.



Source: Authors' elaboration.

1.2. Deforestation emissions accounting method

1.2.1. Bookkeeping models

After revising different methodological approaches to calculate land-use and land-use change emissions, a bookkeeping model approach was chosen, as one of the main advantages is a high traceability, which allows attributing fluxes to specific places, causes, and thus to specific stakeholders. Bookkeeping models make it possible to track carbon stored in vegetation and soils before and after a land-use change event, while excluding the additional ecosystems response to environmental changes (IPCC, 2024).

Therefore, as the present study focuses on land use and land-use change, bookkeeping models are an appropriate choice due to their advantages in Land-use, Land-Use Change, and Forestry (LULUCF) emissions analysis. Additionally, this aligns with the approach taken

by the Global Carbon Budget (GCB), which also relies on bookkeeping models for this specific sector.

Bookkeeping (BK) models estimate emissions from individual land use activities, with a primary focus on agricultural land use changes and the regrowth of secondary forests from abandonment of agriculture. BK models track each land-use and land-cover change (LULCC), allowing fluxes to be presented either as gross emissions and removals or as a net flux. BK approaches combine land cover data with carbon density data for biomass and soil for different ecosystem types. These models track how carbon decays to the atmosphere, or how carbon is removed from the atmosphere following land cover changes (Poulter et al., 2022).

However, it is important to acknowledge the limitations of bookkeeping models. These models depend on predefined parameters and assumptions about carbon density dynamics, typically using exponential, logistic, and linear functions to represent carbon changes. While these functions are based on theoretical frameworks and experimental data, their applicability may vary depending on specific conditions. For the present study, previously established functions will be adopted to ensure consistency and comparability with existing research.

Many studies worldwide have utilized the bookkeeping model approach. Our methodological approach is based on the insights of the following bookkeeping models reviewed for this study: Houghton (Houghton & Castanho, 2023), BLUE (Hansis et al., 2015), LUCE (Qin et al., 2024), and Andersen et al. (2016). All of these models are based on principles established by Houghton et al. (1983), utilize IPCC parameters, and focus on key land-use changes derived from land cover and land cover transition maps. BLUE, LUCE, and Andersen include an assessment at a grid level, which is also a purpose of the present study. In the following section, the equations, assumptions, and adaptations – based on the reference models – will be established to ensure methodological consistency and applicability to the study area.

1.2.2. Carbon Pools

Models typically define the carbon pools considered in the accounting process. In this study, the following carbon pools will be taken into account:

- **Vegetation:** Forest vegetation acts as an emission source when deforested and as a carbon sink when it regrows. Forest vegetation is a key dataset in this study. Both above-ground and below-ground biomass are taken into account. The data sources and dynamics considered are described in the following sections.

- **Soil:** Soil carbon changes following land cover transitions are considered, with emissions occurring when deforestation takes place and carbon sequestration happening during regrowth. Despite uncertainties regarding the soil carbon pool, bookkeeping models usually include soil carbon changes due to land use change as part of the estimations. The estimations are based on soil carbon density data and incorporate temporal response curves to account for changes in soil carbon following land-use transitions. Soil carbon data usually have greater uncertainties than above ground data.

The source for soil properties is provided by FAO's Global Soil Organic Carbon Map (FAO, 2018; GSOCmap v.1.5), the first global map developed through a consultative and participatory process with member countries, coordinated by the Global Soil Partnership. This map can be considered a baseline, as it represents the best available national-level estimate of SOC (FAO & ITPS, 2020). An average value by forest type is calculated from this map and combined with belowground biomass estimates.

- **Atmosphere:** This is the receiving pool of gross emissions, while gross removals extract carbon from the atmosphere, balancing the fluxes of the land-use change process. The atmosphere is assumed to be the final recipient of net emissions resulting from land cover change. Therefore, it is not explicitly discussed in the following sections, as we assume that any net emissions or removals from land cover changes result in an equivalent gain or loss of carbon in the atmosphere.

1.2.3. Land Use and Transitions to be accounted for

The classification of land-use transitions in this study follows approaches from Andersen et al. (2016) and LUCE, considering both anthropogenic and non-anthropogenic changes.


Revising data sources, MapBiomass Bolivia (MapBiomass Bolivia, 2024) arises as the most appropriate source for land cover and land cover transitions maps for Bolivia. This independent source reports the longest available annual data for Bolivia from 1985 to 2023. This source offers an accurate approximation to direct measurements for a vast country like Bolivia, which covers approximately 109,8 million hectares and includes over 50 million hectares of forest.

MapBiomass Bolivia has recently released its Collection 2 of maps, which includes two sets: one for land cover maps and another for land cover transition maps elaborated based on Landsat satellite images with a spatial resolution of 30 meters. The transition maps undergo an additional filtering process to eliminate isolated or edge pixels. Therefore, there may be

minor differences between using land cover and land cover transitions maps (MapBiomass, 2024).

The resulting land cover maps classify the data into six main categories and 19 subcategories. The six primary categories are: Forest, Non-Forest natural formation, Farming, Non-vegetated area, Water body, and Not observed (RAISG, 2024). We adapted this classification following the correspondence table provided in Table 1. An example of the resulting land cover map (for year 2023) is shown on Map 1.

Map 1. Bolivia land cover map, 2023.



Source: Authors' elaboration based on MapBiomass Bolivia (2024). Original land use categories were adapted for this study.

Table 1. Adaptation of the land use types, based on the original MapBiomass classification.

Adapted classification	Original MapBiomass Classification	MapBiomass class codes
Forest	Forest; Open Forest; Flooded Forest	3;4;6
Non-Forest Natural Vegetation	Wetland; Grassland/Herbaceous; Other non-forest natural formation; Shrubland	11;12;13;66
Agriculture	Pasture; Agriculture; Mosaic of uses	15;18;21
Natural non-vegetated area	Beach, dune and sand spot; Rocky outcrop; Salt flat; Other non-vegetated natural area	23;29;61;68
Anthropic infrastructure	Urban infrastructure; Other non-vegetated anthropic area; Mining	24;25;30
Water	River, lake; glacier	33;34


Source: Authors' elaboration based on MapBiomass Bolivia (2024).

Deforestation: Deforestation is taken into account from the pixels changing from forest land cover to other non-forest land cover, annually from 2010 to 2023. The term deforestation typically implies human intervention. According to the Cambridge Dictionary, it is defined as “the cutting down of trees in a large area, or the destruction of forests by people” (Cambridge University Press). In contrast, the FAO (2023) defines deforestation as “the conversion of forest to other land use, regardless of whether it is human-induced or not” emphasizing a change in land use.

Deforestation can be categorized based on its apparent cause and subsequent land use:

- *Clearly Anthropogenic Deforestation:*
 - Conversion of forest to agricultural use, including crops, pasture, and mosaics of uses (as classified by MapBiomass).
 - Conversion of forest to urban areas, mining, or other anthropogenic infrastructure (as classified by MapBiomass).
- *Not Clearly Anthropogenic Deforestation:*
 - Non-agricultural deforestation, where forest loss occurs but there is no visible land use afterward. Although the cause may still be anthropogenic, it is not immediately linked to a subsequent human activity.

Figure 2. Land Use and Transitions considered in this study



Source: Authors' elaboration

To avoid confusion, and given that our data primarily reflect land cover and land cover changes — without reliably distinguishing human causes or capturing actual land use changes not yet reflected in land cover — we use the terms ‘deforestation’ and ‘forest cover loss’ interchangeably throughout this study. Both terms encompass explicitly anthropogenic and not clearly anthropogenic deforestation.

Regenerating Forest (Forest Age Classification): Regenerating Forest is taken into account from pixels changing from any non-forest land cover to forest land cover, annually from 1986 to 2023. Forest age is considered to estimate its carbon absorption. Only the most recent regrowth event for each pixel is taken into account, if the pixel remained classified as forest at least until the beginning of the study period in 2010. Regeneration is categorized based on whether the regrowth occurs after human activities:

- *Anthropogenic Regrowth:* Forest regrowth following agricultural abandonment.
- *Non-Anthropogenic Regrowth:* Forest regrowth from other land uses not directly related to human abandonment.

Throughout this document, ‘regrowth’ encompasses both explicitly anthropogenic and non-anthropogenic regrowth.


A forest type map is used to complement forest information, as MapBiomass Bolivia provides data on land use and transitions but does not include details on forest types. The most recent forest type map was published in 2022 by the Ministry of Environment and Water (MMAyA, 2022). By forest extension the main type in Bolivia is the Amazon forest followed by Chaqueño and Chiquitano forest. The 9 major forest types identified in the data source are used for the present analysis.

Since the available map identifies areas covered by the different forest types only for the year 2022, we created a “forest type zone” map by combining the 2022 forest type data with an ecoregion map (Ibisch & Merida, 2003). By integrating the Forest Type dataset (MMAyA, 2022) with the Ecoregion dataset (Ibisch & Mérida, 2003), we produced a new dataset comprising the nine forest categories defined by MMAyA, applied across the entire country. This represents a potential forest type map for Bolivia, assuming full forest cover.

This adjustment is essential because our land use and transition analysis covers both current forest areas and historical forest changes, while the 2022 forest type map only reflects areas that remained forested as of that year, excluding areas deforested earlier. In the deforestation analysis, it allows us to identify the original forest type associated with each land cover change. In the fire emissions analysis, the forest type map is used as a

reference to determine the ecozone, even in areas that are currently non-forest or have never been forested, helping to contextualize fire dynamics across different ecological regions.

Map 2. Current and potential forest type in Bolivia.



Note: Left: Original forest type map of 2022 (MMAyA, 2022). Right: Potential forest type zone. Source: Authors' elaboration.

While deforestation emissions are calculated at a 30x30 meter resolution, results are presented at a 100x100 meter resolution. The rescaling method used for the MapBiomass and forest type data is based on the 'nearest neighbor' approach, which is the recommended method for rescaling categorical (class) data.

1.2.4. Assumptions for deforestation estimations

To estimate CO₂ emissions from deforestation with models, it is necessary to make some assumptions. The most important ones are presented below:

- **Aboveground biomass is almost completely removed**

The assumption that aboveground biomass is almost completely removed after deforestation to agricultural use is widely applied in bookkeeping models. As Hansis et al. (2015) states: "*For clearing, it is assumed that the respective biomass of the source type is completely removed, with none leftover in the target cover type.*" Similarly, Andersen et al. (2016) assumes that after a land-use change to agriculture, aboveground biomass is reduced to the level of crops or pastures biomass. This assumption is kept for the present

study. The year of a deforestation event, the baseline biomass changes to the destination land covers, which are explained below.

- Deforestation to agriculture or pasture: The biomass content is adjusted to the default values for these land uses, following IPCC (2006) guidelines.
- Deforestation to urban areas: Aboveground biomass is assumed to be completely removed (zero biomass).
- Deforestation with no apparent land use (non-use deforestation): The remaining biomass is assumed to correspond to the average natural non-forest vegetation biomass, estimated using Santoro & Cartus (2024) biomass maps overlaid with natural non-forest vegetation cover from MapBiomass, by forest type zones.
- **Belowground carbon is assumed to decompose gradually.**

Belowground carbon, which includes soil carbon and belowground biomass, is assumed to decompose gradually after deforestation following a linear function over 20 years, ultimately resulting in a 25% loss of its initial carbon content.

- Soil carbon averages are obtained from soil maps. Values are estimated as an average by forest type to account for spatial variability.
- Belowground biomass is estimated using average carbon content in intact forests, stratified by forest type. The IPCC (2006) root-to-shoot ratios are applied to estimate belowground biomass.
- As assumed in Andersen et al. (2016) based on literature, we assume no carbon emissions from soils if there is no apparent agricultural use after deforestation, when forests turn to natural non-forest cover.

1.2.5. Equations and Parameters

- **Establish biomass baseline by forest type**

Intact forest since 1985 was identified using land cover and transition data from MapBiomass Bolivia (2024). After extracting this area (in pixels), this undisturbed forest is utilized to extract the aboveground biomass from the biomass map provided by Santoro & Cartus (2024) available at ESA Biomass Climate Change Initiative (CCI Biomass). This source makes available global aboveground biomass maps for the year 2010 and from 2015 annually till 2021 with a 100 meter pixel resolution. Biomass estimates are important for assessing the

carbon removal (or emission) capacity of different types of ecosystems, as it directly correlates to an ecosystem's ability to sequester or release carbon.

As with other bookkeeping models, this study uses a single baseline biomass dataset from the year 2010 to estimate potential carbon emissions or absorptions based on the average biomass of each forest type. Average biomass in intact forests was also calculated for the years 2015 to 2021, yielding similar results and thereby ensuring consistency in biomass estimates for intact forest areas. While we use average estimates of biomass for each type of forest for our central emissions estimates, it is important to keep in mind that deforestation and forest fires may not target average forest, but rather less dense forest. This possibility will be explored in the sensitivity analysis.

Belowground biomass is estimated by applying the R factor (Ratio of Belowground Biomass to Aboveground Biomass) (from table 4.4 in Volume IV, Section 4, (IPCC, 2019, p. 18)), which represents the relationship between below-ground and above-ground biomass, as direct measurements or estimations are not available.

Table 2. Aboveground and belowground biomass data by forest types (tons of biomass/ha).

Forest type	Aboveground Biomass				Belowground Biomass	
	Average biomass (t/ha)	Percentile 75th (t/ha)	Percentile 50th (t/ha)	Percentile 25th (t/ha)	Ratio of belowground to aboveground biomass	Average belowground biomass (t/ha)
Amazónico	224	263	221	184	0,221	49
Yungas	189	229	186	145	0,283	54
Chiquitano	152	181	157	127	0,284	43
Tucumano – Boliviano	144	201	152	101	0,283	41
Llanuras Inundables	143	172	129	77	0,221	32
Pantanal	107	132	115	76	0,285	31
Seco Interandino	65	102	57	22	0,348	23
Chaqueño	54	69	43	27	0,334	18
Andino	50	82	35	7	0,348	17

Note: The table presents values in tons of biomass; carbon content is assumed to be half of the biomass, as is commonly established. Source: Authors' elaboration based on Forest map 2022 (MMAyA, 2022), Land cover and transitions maps (MapBiomass Bolivia, 2024), Biomass map (Santoro & Cartus, 2024) and IPCC default values for ratios of belowground biomass.

- **Forest Regrowth curves**

The underlying principle of the following equation, as seen in the bookkeeping models BLUE and LUCE, is that carbon reservoirs do not remain disturbed indefinitely. Instead, they follow

a predictable trajectory toward a new stable state determined by post-disturbance land use.

o **Aboveground biomass**

In the case of regenerating forests, carbon content increases rapidly in the early years, then slows as the forest matures, eventually reaching a stable carbon level typical of mature forests. This predictable pattern allows for simplified modelling without significant loss of accuracy. Since various models adopt similar functional approaches, we apply the function proposed by Andersen et al. (2016), as shown below:

$$CAR_{v,a} = \frac{CAP_v}{1 + e^{\alpha_v - \beta_v a}}$$

Where $CAR_{v,a}$ represents the amount of carbon during regeneration at age a for a forest type v and CAP_v is the amount of carbon of a mature forest of the same type v . The parameters α_v and β_v determine the exact shape of the logistic function which vary across the nine forest types as shown in Figure 3.


Based on the parameters established by Houghton and Hackler (2001), it is estimated that tropical rainforests in Bolivia take 40 years to regenerate and reach the carbon content of intact forests, while dry or seasonal forests require 35 years to recover their original carbon content. Based on the Bolivian forest types, four of the nine forest types can be considered dry forest (*Andino*, *Chaqueño*, *Seco Interandino* and *Tucumano Boliviano*) while the other five are most likely tropical rainforest (*Amazónico*, *Chiquitano*, *Llanuras Inundables*, *Pantanal* and *Yungas*).

To calculate the parameters α_v and β_v , two assumptions are made. First, different initial carbon content values are assumed at the beginning of regeneration. Based on the fact that most deforestation occurs to make way for agricultural uses, the regeneration curves assume carbon recovery begins from the minimum aboveground carbon content typical of agricultural uses, specific to each forest type. These values range from 2 tC/ha for most forest type zones to 10 tC/ha in the Amazonian and Yungas forest type zones, according to IPCC (2019).

Second, forest regeneration is assumed to take between 35 and 40 years to reach 99% of its maximum carbon content. The average maximum carbon content by forest type is the one estimated as the baseline biomass since it represents average biomass content in old forest and is the value assumed for variable CAP_v , which varies across the nine forest types, ranging from 25 tC/ha in the Andean forest type to 112 tC/ha in the Amazonian forest type. Taking into

account the time to recover and the maximum reached carbon content, nine separate logistic functions were estimated, resulting in a unique regrowth curve for each forest type.

Figure 3. Evolution of aboveground carbon content in regrowing forests, by forest type.



Source: Authors' elaboration.

o **Belowground biomass**

As noted by Andersen (2016), various literature reviews indicate that soil carbon is resilient to aboveground disturbances, remaining relatively stable even in degraded forests or areas converted to pasture. Based on the assumptions of Houghton and Hackler (2001), the model assumes that belowground carbon regenerates linearly, following a curve with half the slope of the belowground carbon decomposition curve.

1.3. Assessment of burned area and fire emissions data

Fire emissions are not always included in AFOLU emissions estimations. While some estimations include emissions from burned peatlands, fire emissions have often been calculated separately from LULUCF emissions. Studies like the Global Carbon Budget have only recently begun incorporating fires into their estimations and separated from LULUCF estimations.

Some burned forest may be completely lost and thus accounted for as deforestation by transition maps. However, other burned forest may remain as forest, but degraded. In these cases, transitions maps will not identify them as forest loss. Fires and fire emissions require separate analysis to understand their impact on forest loss, as well as their dynamics with land cover transitions, forest degradation, and regeneration.

Besides the difficulties in distinguishing human-caused fires from natural ones (GCB, 2025), other difficulties to estimate fire emissions are the variations in fire intensity, and the complexity of post-fire ecosystem recovery. On-the-ground observations are often impractical, making satellites the primary tool for monitoring. However, remote sensing has limitations, such as misidentifying fire signals and struggling to detect small fires. Ground-based data is often inconsistent and difficult to access, complicating emissions model validation. Estimating emissions remains complex, as it depends on factors such as vegetation type and fuel composition of vegetation, fire intensity, and landscape features.

These challenges make it difficult to perform primary estimations of fire emissions. Therefore, to incorporate fire emissions into the analysis, existing fire emission datasets will be used to assess their impact and relevance in Bolivia. This analysis requires two key data sources and their analysis:

- **Burned Area** which will enable the assessment of land cover affected by fires and other spatial analyses.
- **Fire emissions** to analyse and utilize global fire emission estimation data to assess the impact, magnitude, and other relevant characteristics of fires in Bolivia. Integrating these estimates with deforestation-related emissions will help ensure the avoidance of double counting.

Concerning burned area, given the available data and the need to analyse the period from 2010 to 2023, it seems reasonable to use MODIS data on burned areas, specifically the MCD64A1 Version 6.1 product. This product is derived from MODIS satellite images and provides burned area maps based on calculations from its sensors (Giglio et al., 2019).

For fire emissions, the Global Fires Emissions Database (GFED) (Van Wees et al., 2022) is used. The GFED is based on a global fire emissions model at a 500 m spatial resolution, integrating multiple remote sensing datasets to estimate biomass burning fuel consumption and emissions. The GFED framework estimates emissions from satellite-based data on vegetation cover, productivity, and burned areas. From the information available, the national territory of Bolivia is extracted for the period 2010–2022, since there's no data available for 2023. The main concern regarding this segment of the analysis is the potential for double counting emissions in areas where both fire and deforestation have occurred, as deforestation emissions are already included in the bookkeeping calculations. To avoid this overlap and obtain an approximation of degradation emissions we can use the following assumption:


Degradation emissions = Total fire emissions – deforestation emissions within burned areas

Where:

- **Total fire emissions** include emissions from deforestation (meaning total forest cover loss), forest degradation, and non-forest vegetation degradation. These estimates are based on data from the GFED.
- **Deforestation emissions** refer to emissions resulting from the complete conversion of forest cover to other land cover types. These are primarily estimated using the bookkeeping model, incorporating spatial and temporal overlap with burned areas. Deforestation emissions occur both within and outside burned areas. Only deforestation emissions inside burned areas should be considered to avoid double-counting.
- **Degradation emissions** represent estimated emissions from fires affecting both forest and non-forest vegetation. The resulting degradation emissions include both forest and non-forest vegetation degradation. In the case of forests, this includes areas that remain classified as forest in land cover maps despite being affected by fire.

Both deforestation and forest degradation contribute to greenhouse gas emissions. Fires may destroy the forest entirely (causing deforestation) in some cases, while in other cases, they may leave a degraded forest with lower quality and productivity (Lanly, 2003). The overlaps between forest, burned areas, degradation and deforestation is presented on Figure 4.

Figure 4. Forest, burned area, degradation and deforestation overlaps



Source: Authors' elaboration.

Although many gases are emitted from fires, numerous sources agree that the primary greenhouse gas (GHG) released is carbon dioxide (CO₂), with methane (CH₄), nitrous oxide (N₂O) and others being minor components (Sims et al., 2024). These emissions are generally estimated based on the amount of biomass burned. For the present analysis, all biomass burned will be considered as CO₂ emissions, since CO₂ accounts for more than 90% of fire-related emissions. This approach also facilitates comparison with deforestation emissions.

It is important to take into account that fire CO₂ emissions from the Global Fires Emissions Database represent gross carbon fluxes, while carbon recovery post-fire is not fully accounted for (Friedlingstein et al., 2023).

Note that the biomass data used in GFED differs from the dataset employed in this study, introducing an additional source of uncertainty in this specific calculation. Therefore, these results are included in the discussion section rather than in the main results section, as they represent preliminary estimates that require further analysis to reduce uncertainty.

All fire-related data is available at a pixel resolution of 500 meters. Since the fire analysis was conducted separately, its results are also at a 500-meter resolution. When this data is compared with deforestation data – which is mostly analyzed at aggregated scales such as national, municipal or forest type zones – the aggregation ensures that the results remain comparable.

2. Results

An Excel workbook with all the calculations and results at the municipal level accompanies this article, as do raster maps of net carbon emissions from deforestation at the 100x100m resolution and total fire emissions at the 500x500m resolution. All files can be freely downloaded from the following folder:


<https://drive.google.com/drive/u/0/folders/1BoXRaWo55GiDYfh8HUhlWp9FIKfHh-x->

2.1. Carbon emissions from land-use change – Deforestation

2.1.1. Gross emissions

To estimate gross emissions, from both aboveground and belowground carbon content (including belowground biomass and soil carbon), forest loss since 2010 was taken into account. The deforested surface tends to increase over time, and it is mostly located in the department of Santa Cruz. Figure 5 shows the annual forest loss over 2010–2023.

Figure 5. Annual Forest loss in Bolivia, 2010 – 2023 (ha/year).




Source: Authors' elaboration based on MapBiomass Bolivia (2024).

Emissions occur where deforestation takes place; however, the amount of emissions per hectare varies depending on the forest type. Over the entire analysis period, the highest emissions per hectare are found in the northern regions of Bolivia, even though the total amount of deforestation is greater in the eastern part of the country. Similarly, within the department of Santa Cruz – which has the highest total deforestation – emissions per hectare are greater in the north, where forest types are predominantly Amazónico and Chiquitano. In contrast, the southern part of the department, dominated by Chaco and Pantanal forests, shows lower emissions per hectare.

For reporting and comparability purposes, the assumption is that all carbon emissions are released as CO₂. Carbon (C) emissions are converted to carbon dioxide (CO₂) using a conversion factor of 3,664. Map 3 shows spatial distribution of gross emissions while Table 3 presents results of cumulative gross emissions and deforested area during the study period by forest type.

Map 3. Total gross emissions from forest loss in Bolivia 2010–2023



Source: Authors’ elaboration.

Table 3. Total deforestation and gross emissions by forest type.


Forest type	Deforestation, 2010 – 2023 (ha)	Total Gross emissions, 2010 – 2023 (tCO ₂)
Amazónico	1.390.986	515.836.471
Chiquitano	1.407.882	389.084.009
Chaqueño	1.370.162	139.716.441
Llanuras Inundables	451.175	112.531.968
Yungas	214.549	70.962.057
Tucumano – Boliviano	126.311	34.225.703
Pantanal	151.439	29.497.381
Seco Interandino	32.216	3.873.437
Andino	3.750	342.157
Bolivia	5.148.469	1.296.069.625

Note: Deforestation includes the total area that has lost forest between 2010 and 2023. Source: Authors’ elaboration.

2.1.2. Gross removals

To estimate gross removals (Map 4 and Table 4), regrowing forests established since 1985 and still standing at the beginning of our analysis period were considered, accounting for their carbon removals starting from 2010. Based on forest regrowth curves, younger forest removes higher carbon quantities than older forest. The capacity to remove carbon also depends on the forest type.

Map 4. Total gross removals from forest regrowth in Bolivia 2010–2023.



Source: Authors' elaboration.

Table 4. Total forest regrowth and gross removals by forest type.


Forest type	Regrowing forest by 2023 (ha)	Total Gross removals, 2010 – 2023 (tCO ₂)
Amazónico	1.467.242	185.808.365
Chiquitano	587.997	38.694.366
Llanuras Inundables	608.884	33.262.695
Yungas	279.453	30.215.575
Chaqueño	469.506	16.844.942
Tucumano – Boliviano	122.778	10.901.629
Pantanal	95.455	4.259.554
Seco Interandino	38.866	1.441.251
Andino	5.017	158.205
Bolivia	3.675.199	321.586.581

Note: The regrowing forest by 2023 includes forest that have been regenerating since the first transition map, representing forest aged between 1 and 37 years. Source: Authors' elaboration.

2.1.3. Net fluxes

The difference between gross emissions from deforestation and gross removals from forest regrowth are the net emissions. Both gross and net emissions reveal an increasing trend (Figure 6 and Table 5).

Figure 6. Annual gross and net fluxes in tCO₂/year, 2010 – 2023.



Source: Authors' elaboration.


Table 5. Annual gross and net fluxes in tCO₂, 2010 – 2023.

Year	Gross emissions	Gross removals	Net emissions
2010	70.298.334	15.596.420	54.701.914
2011	114.008.338	16.598.778	97.409.559
2012	88.444.942	17.325.381	71.119.561
2013	61.195.298	18.592.989	42.602.309
2014	79.514.407	19.401.399	60.113.008
2015	63.837.235	20.807.307	43.029.928
2016	84.264.743	21.862.016	62.402.727
2017	87.342.563	23.124.118	64.218.445
2018	87.690.467	24.298.907	63.391.560
2019	106.894.064	25.357.840	81.536.224
2020	91.211.892	26.950.907	64.260.985
2021	127.589.997	28.961.165	98.628.832
2022	103.342.167	31.076.186	72.265.981
2023	130.435.179	31.633.169	98.802.010
Total 2010 – 2023	1.296.069.625	321.586.581	974.483.044

Source: Authors' elaboration.

The net emissions map can be obtained spatially by overlapping maps of gross emissions and gross removals. The next map shows the distribution of net emissions.

Map 5. Net emissions in Bolivia 2010–2023.



Source: Authors' elaboration.

Geographical differences can be explained by referring to two earlier maps in this document: the 2023 land cover map (Map 1) and the 2022 potential forest type map (Map 2). These maps show that the areas with the highest net emissions largely coincide with agricultural land cover in 2023, indicating that this expansion of agriculture resulted from forest loss. Additionally, the potential forest type map reveals that the southern part of Santa Cruz — dominated by Chaco forest — shows lower net emissions, as this forest type has lower biomass density (see Table 2). In contrast, the northern areas of Santa Cruz, characterized by Chiquitano and Amazon forests, exhibit higher net emissions per hectare due to their greater biomass content.

It is interesting to analyse cases by forest type separately. For example, Amazon forest has the highest gross emissions and highest gross removals ranking second in the net emissions, while Chiquitano forest has lower removals reaching the first place in net emissions (Table 6).

Table 6. Cumulative gross and net fluxes by forest type in tCO₂, total of the period 2010 – 2023

Forest type	Gross emissions	Gross removals	Net emissions	% of total net emissions
Chiquitano	389.084.009	38.694.366	350.389.643	36%
Amazónico	515.836.471	185.808.365	330.028.107	34%
Chaqueño	139.716.441	16.844.942	122.871.499	13%
Llanuras Inundables	112.531.968	33.262.695	79.269.273	8%
Yungas	70.962.057	30.215.575	40.746.482	4%
Pantanal	29.497.381	4.259.554	25.237.827	3%
Tucumano – Boliviano	34.225.703	10.901.629	23.324.074	2%
Seco Interandino	3.873.437	1.441.251	2.432.186	0%
Andino	342.157	158.205	183.952	0%
Bolivia	1.296.069.625	321.586.581	974.483.044	100%

Source: Authors' elaboration.

Table 7 shows results relative to population and GDP, revealing an average annual emission of 6,6 tCO₂ per capita, much higher than the global average of emissions from land-use change of about 1,4 tCO₂ per capita (Global Carbon Atlas, 2023).

Table 7. Annual net fluxes per capita and per GDP unit, 2010 – 2023.


Year	Net emissions (tCO ₂)	Net emissions per capita (tCO ₂ /person)	Net emissions per GDP unit (kgCO ₂ /Bs)
2010	54.701.914	5,6	1,7
2011	97.409.559	9,8	2,8
2012	71.119.561	7,1	2,0
2013	42.602.309	4,2	1,1
2014	60.113.008	5,9	1,5
2015	43.029.928	4,1	1,0
2016	62.402.727	6,0	1,4
2017	64.218.445	6,1	1,4
2018	63.391.560	5,9	1,3
2019	81.536.224	7,6	1,7
2020	64.260.985	5,9	1,4
2021	98.628.832	9,0	2,1
2022	72.265.981	6,5	1,5
2023	98.802.010	8,8	1,9
Average		6,6	1,6

Note: Bs. is the abbreviation for the local currency, Bolivianos. At the official exchange rate, 1 Bs. ≈ USD 0,14, while in the informal market, 1 Bs. ≈ USD 0,07. Source: Authors' elaboration.

Forest cover loss can result from both human activities and natural factors. However, even when deforestation does not lead to an immediately clear human land use, it cannot be definitively ruled out as anthropogenic. In contrast, deforested areas where agriculture or infrastructure development is observed can be more confidently attributed to human-

driven deforestation, and thus anthropogenic emissions. To ensure a comprehensive analysis, forest cover gains from agricultural abandonment were also considered as part of human-induced land-use changes. The results of this disaggregation are presented in Figure 7 and Table 8.

Figure 7. Anthropogenic and total annual Net fluxes (tCO₂), 2010 – 2023.



Source: Authors' elaboration.

Table 8. Annual gross and net anthropogenic fluxes (tCO₂), 2010 – 2023.

Year	Gross emissions	Gross removals	Net emissions	Total net emissions from clearly anthropogenic emissions (%)
2010	41.925.118	5.127.005	36.798.112	67%
2011	60.611.036	5.512.947	55.098.090	57%
2012	48.425.671	5.902.678	42.522.993	60%
2013	38.533.133	6.509.418	32.023.715	75%
2014	41.385.545	6.843.278	34.542.267	57%
2015	41.789.444	7.342.381	34.447.062	80%
2016	64.236.451	7.735.225	56.501.227	91%
2017	67.530.568	8.453.428	59.077.140	92%
2018	65.522.805	9.017.396	56.505.409	89%
2019	70.073.983	9.530.493	60.543.490	74%
2020	63.176.549	10.142.618	53.033.931	83%
2021	92.281.776	10.632.623	81.649.153	83%
2022	58.011.169	10.968.795	47.042.374	65%
2023	109.928.948	12.333.818	97.595.131	99%
Total 2010 – 2023	863.432.196	116.052.103	747.380.094	77%


Source: Authors' elaboration.

2.2. Burned area and fire emissions

2.2.1. Burned area

As previously mentioned, the primary source for burned area data in this study is the MODIS burned area product. According to this dataset, burned area trends do not follow a stable pattern. However, in recent years, there has been an overall increase in the total burned area compared to the period before 2018. An exception is 2010, which recorded an unusually high value due to a prolonged dry season and limited government response capacity at the time, allowing traditional land-clearing fires for agricultural expansion to spread over much larger areas than in other years. Figure 8 illustrates the total annual burned area in Bolivia from 2010 to 2023 according to MODIS data.

Figure 8. Total annual burned area in Bolivia, in hectares, 2010–2023.



Source: Authors' elaboration based on MODIS (Giglio et al., 2019).

Figure 9. Composition of land-cover types in annual burned areas in Bolivia, 2010–2023.




Source: Authors' elaboration based on MapBiomass Bolivia (2024) and MODIS (Giglio et al., 2019)

Significant differences emerge when the data is disaggregated by land cover type. Figure 9 depicts this variation across different land cover categories. Natural non-forest vegetation is the main land cover within burned areas, followed by forest with a worrying 40% of the land cover affected by fire in 2019 and it exceeds 20% since then.

Deforestation areas reach a maximum of about 500 thousand hectares annually and the average is about 360 thousand hectares per year. On the other hand, burned areas reached a total of more than 9 million hectares in 2010 and the minimum per year is over 1 million hectares. Forest within burned areas reached 3 million hectares in 2010 and 2 million hectares in 2019. Given these data we can conclude that forests are importantly affected by fires but only a small proportion of fires result in complete forest cover loss (deforestation).

Another important question is: How much of total deforestation occurs within burned areas? It is evident that deforestation is not a high proportion of burned areas, but burned areas represent a quite high proportion of the total deforested areas. On average, 40% of deforestation occurs within burned areas during the period of analysis (Figure 10).

Figure 10. Forest loss in burned areas in Bolivia in hectares, 2010–2023.



Source: Authors' elaboration based on MapBiomass Bolivia (2024) and MODIS (Giglio et al., 2019).


The analysis of burned areas suggests that fire emissions are primarily associated with the degradation of forest and non-forest vegetation, as the extent of deforestation within burned areas is smaller than that of burned forest and total burned areas.

2.2.2. Fire emissions estimates 2010 – 2022

As previously mentioned, the data on fire emissions – expressed in terms of biomass burned and converted to carbon emissions – used in this analysis is sourced from the Global Fire Emissions Database (GFED). For consistency and ease of comparison with CO₂ emissions from deforestation, we focus on CO₂ emissions (Van Wees et al., 2022). Spatial distribution of biomass burned offered by GFED is presented on map 6.

To identify the ecosystems most affected by fires, the “forest type zone” map is used as a reference to determine ecozones. This approach helps contextualize fire dynamics across broader ecological regions, recognizing that these zones include a variety of land cover types – not just forests. Based on this classification, and considering all areas affected by fire at least once between 2010 and 2022, the *Llanura Inundable* (flooded savanna) in the department of Beni stands out as the most impacted region. These savannas are predominantly wet for most of the year, but during the dry season (approximately June to October), they dry out considerably – coinciding with the peak fire season, often driven by agricultural practices.

However, because this area is primarily covered by non-forest vegetation, its fire-related emissions are relatively low. In contrast, the Chiquitano and Amazonian forest zones, where forests are the dominant land cover type, experience significantly higher fire emissions. This difference is reflected in Table 9.



Source: Authors' elaboration based on GFED.

Table 9. Total burned area and total fire emissions by forest type zones, 2010 – 2022.

Forest type zone	Total burned area 2010 – 2022 (ha)	Total fire emissions 2010 – 2022 (tCO ₂)
Chiquitano	3.615.175	528.611.000
Amazónico	3.234.275	386.069.985
Llanuras Inundables	8.682.750	333.915.528
Chaqueño	1.936.375	236.894.687
Pantanal	2.058.250	185.871.735
Tucumano – Boliviano	205.700	14.506.536
Yungas	112.875	8.338.657
Seco Interandino	35.650	856.809
Andino	95.800	796.610

Note: “Total burned area” refers to the cumulative extent of land affected by fire during the study period, regardless of the number of times it burned. For example, if a surface burned multiple times, it is counted only once in this table. This differs from fire emission estimates, where emissions are accounted for each year a fire is recorded. Source: Authors’ elaboration.

2.2.3. Fire emissions from forest degradation

Fire emissions estimates account for all burned vegetation emissions, including both forest and non-forest biomass. Also, these estimates encompass forest loss emissions as well as emissions from burned forests that remain as forest.

To estimate the vegetation (forest and non-forest) degradation component, the approach used is to subtract deforestation-related emissions within burned areas from the total fire emissions estimates. Since previous results indicate that burned areas are significantly larger than deforested areas, and that less than half of the deforested areas are burned, the contribution of forest loss emissions within total fire emissions is expected to be relatively low. The results in Table 10 are consistent with the expected results.

Table 11 shows that most of the areas impacted by fires correspond to natural non-forest vegetation. However, we cannot attribute the majority of emissions to this type of vegetation degradation, since forests store significantly larger quantities of biomass. This is also evident in Map 6, where the department of Beni — with a large extent of natural non-forest vegetation — shows a wide area affected by fire emissions. In contrast, the department of Santa Cruz — where a greater extent of burned forest is found — exhibits higher fire emissions in terms of tons of carbon per hectare.

Table 10. Fire emissions from degradation (tCO₂), 2010 – 2022.

Year	Total Fire emissions	Deforestation emissions within fire emissions area	Estimated emissions by vegetation degradation (forest and non-forest)
2010	327.003.752	6.321.279	320.682.473
2011	95.939.847	10.206.842	85.733.005
2012	63.349.127	7.841.899	55.507.228
2013	37.932.645	5.323.794	32.608.851
2014	32.535.906	6.901.800	25.634.106
2015	63.333.713	5.476.081	57.857.632
2016	123.506.171	7.258.626	116.247.545
2017	83.220.377	7.466.187	75.754.190
2018	74.671.144	7.417.110	67.254.034
2019	254.211.134	9.046.468	245.164.666
2020	173.223.478	7.544.756	165.678.722
2021	188.544.798	10.770.227	177.774.571
2022	182.655.519	8.494.051	174.161.468

Source: Authors' elaboration.

Table 11. Land cover composition in burned area in 2023 by forest type zone.

Forest type zone	Natural non-forest vegetation	Forest	Pasture	Agriculture	Mosaic of uses	Non-vegetated area	Total
Llanuras Inundables	52,46%	5,42%	6,54%	0,17%	0,06%	0,43%	65,08%
Amazónico	6,56%	14,24%	0,95%	1,47%	0,29%	0,21%	23,72%
Pantanal	3,77%	0,27%	0,09%	0,00%	0,02%	0,00%	4,16%
Chiquitano	0,32%	1,39%	0,50%	0,62%	0,43%	0,01%	3,28%
Chaqueño	0,66%	1,25%	0,21%	0,59%	0,06%	0,01%	2,79%
Yungas	0,19%	0,48%	0,00%	0,02%	0,01%	0,01%	0,71%
Andino	0,13%	0,01%	0,00%	0,00%	0,00%	0,03%	0,17%
Seco Interandino	0,04%	0,01%	0,00%	0,00%	0,00%	0,00%	0,05%
Tucumano – Boliviano	0,02%	0,01%	0,00%	0,00%	0,01%	0,00%	0,04%
Total	64,15%	23,08%	8,29%	2,88%	0,89%	0,71%	100,00%

Note: This table does not reflect forest cover loss; it only shows land cover types as classified in the year 2023 within burned areas. Source: Authors' elaboration.

To separate forest degradation from non-forest vegetation degradation is not an easy task. For example, if a fire emission pixel shows a total of 100 tons of carbon (in a 500m pixel), the pixel could be composed half of non-forest and half of forest. However, those 100 tons of carbon emissions will not be evenly split between the two. The majority of emissions are likely to come from the forest portion due to its higher biomass. Depending on the forest type zone, the median carbon content in natural non-forest vegetation ranges from nearly 0 tC/ha in


the *Andino* forest type zone to 20 tC/ha in the *Amazónico* forest type zone. Moreover, the differences between the average and median values are much greater in natural non-forest vegetation than in forests. Therefore, this separation needs further analysis.

Another important consideration is that the estimated degradation emissions represent gross emissions. While fire-induced carbon losses may be partially offset by post-fire vegetation regeneration, this process is neither consistent nor easily predictable. As discussed below, evidence shows that regeneration does not always occur following fire events, and does not follow the same behaviour everywhere.

For example, a study conducted by Maillard (2023) aimed to estimate post-fire regeneration trends in Bolivian ecosystems using the Normalized Difference Vegetation Index (NDVI), a widely used indicator for assessing vegetation dynamics derived from satellite imagery. One key finding of the study was the recurrence of fires, meaning that some areas burn repeatedly over different years. The research also highlighted that different ecosystems respond differently to fire. For instance, the Chiquitano and Chaco ecoregions are better adapted to recover from fires, as many of their plant species are resilient to dry conditions and occasional natural fires. However, even in these fire-adapted regions, post-fire regeneration does not always occur. Maillard (2023) found that 54% of the burned areas in Bolivia showed a significant increase in NDVI, indicating vegetation regeneration. Meanwhile, 30% of the areas exhibited mixed trends — both increasing and decreasing NDVI values — but these trends were not statistically significant. In contrast, 16% of the burned areas displayed a significant decreasing NDVI trend, suggesting ongoing degradation. Nearly half of the areas showing signs of regeneration were located in savannas, particularly in the department of Beni, where forest cover is minimal and the landscape is predominantly composed of natural non-forest vegetation.

Based on MODIS data from 2010 to 2023, we estimated the recurrence of fires across the same areas. The analysis shows that most burned areas were affected by fire in more than one year, either in consecutive or non-consecutive years. As shown in the following graph, 44% of the total burned area burned only once during the 2010–2023 period, while the remaining 56% burned in two or more years.


Figure 11. Recurrency of burned areas, 2010–2023.



Source: Authors' elaboration based on MODIS (Giglio et al, 2019).

In addition, we estimated biomass changes looking forward to assess emissions from forest degradation caused by fire. The results show that biomass in unburned areas remained generally stable. In contrast, biomass in burned areas exhibited fluctuations with a slight decreasing trend, while areas that burned more than once showed more pronounced fluctuations and a clearer downward trend. Figure 12 shows the general trends.

Figure 12. Average Forest biomass within burned areas, 2015–2021.



Source: Authors' elaboration based on MODIS (Giglio et al, 2019) and Santoro & Cartus (2024).


2.3. National results

Figure 13 summarizes the average annual carbon flows between different pools for the period 2010–2023. The largest flow – 126 million tons of CO₂ per year – comes from fires that degrade forests without completely removing forest cover and to a lesser extent, from the degradation of natural non-forest vegetation. This is particularly concerning, as these emissions occur without any associated land-use change or productive land use.

The second largest flow—62 Mt CO₂ per year—is from deforestation driven by conversion of forest to agricultural land, while the third most important flow is from forest to natural non-

forest (31 Mt CO₂ per year). The latter changes are also problematic, as they appear to generate no clear economic benefit while contributing to forest loss. However, there are also significant changes from natural non-forest to forest, highlighting the dynamic and reversible nature of some land cover changes. This environmental fluidity adds complexity to estimating carbon emissions and removals.

Figure 13. Average annual carbon flows from land use change and fires in Bolivia, 2010–2023.



Source: Authors' elaboration.

2.4. Subnational level results

2.4.1. Departmental level


While total emissions from deforestation are highest in Santa Cruz, emissions per capita and per GDP unit are higher in Beni and Pando (see Table 12; see Map 1 for departmental boundaries). Fire emissions follow a similar pattern, as Santa Cruz is the department with highest emissions followed by Beni, while the remaining departments have experienced very low fire emissions most years.

Table 12: Net emissions by deforestation by department, 2010 – 2023.

Department	Total Net Emissions (tCO ₂ 2010–2023)	Average tCO ₂ /year	Average tCO ₂ /year/person	Average kgCO ₂ /year/Bs.
Santa Cruz	659.093.830	47.078.131	16,6	3,4
Beni	173.726.700	12.409.050	27,9	8,6
La Paz	51.700.108	3.692.865	1,3	0,3
Pando	34.346.552	2.453.325	20,8	6,6
Cochabamba	30.172.216	2.155.158	1,2	0,3
Tarija	14.728.477	1.052.034	2,1	0,3
Chuquisaca	10.676.571	762.612	1,3	0,4
Potosi	38.439	2.746	0,0	0,0
Oruro	151	11	0,0	0,0
Bolivia	974.483.044	69.605.932	6,6	1,6


Source: Authors' elaboration.

Figure 14. Net land-cover change emissions, by department, 2010–2022 (tCO₂/year).



Source: Authors' elaboration.

Figure 15. Total fire emissions (tCO₂/year), by department, 2010–2022.




Source: Authors' elaboration.

2.4.2. Municipal level

In Bolivia there are 343 municipalities. Only 7 of them are net carbon sinks during the period of analysis with a maximum annual average of – 65 tCO₂ net emissions in the municipality of Icla, department of Chuquisaca. Most municipalities especially in the southwest has almost zero emissions since there is almost no forest cover. While the east concentrates the municipalities with higher emissions.

Map 7. Average Net emissions from deforestation (tCO₂/year), 2010–2023.



Source: Authors' elaboration.

To identify the municipalities with the highest net emissions from land-cover change, we used a combination of three key criteria: 1) Highest absolute emissions; 2) Highest emissions per capita; and 3) Highest emissions as a percentage of municipal biomass stock. The top 25 municipalities identified by each criterion are presented in Table 13.


Map 8 shows the top 25 municipalities by each criterion, and the intersection between criteria. In black, we see seven municipalities that rank within the top 25 according to all three emissions criteria. All are located in the department of Santa Cruz. Almost all of the municipalities losing a high percentage of their carbon stock are also located in this department, whereas municipalities with high per capita emissions, are found mainly in Pando and Beni. The black municipalities are clearly the most concerning.

Table 13. Top 25 municipalities – by different measures of net emissions from deforestation, 2010–2023

Absolute net emissions			Percentage of municipality's carbon stock		Emissions per capita	
Municipality	Average tCO ₂ /year	% of country emissions	Municipality	% emissions/carbon stock	Municipality	Average tCO ₂ /person/year
San Ignacio de Velasco	8.327.712	12%	Colcapirhua	4,18%	El Carmen Rivero Tórrez	317
El Puente	3.657.417	5%	San Julián	4,09%	Puerto Siles	243
San José de Chiquitos	3.381.242	5%	Okinawa Uno	3,00%	El Puente	223
Concepción	3.347.241	5%	Cuatro Cañadas	2,23%	San Rafael	222
Ascención de Guarayos	2.747.962	4%	Santa Rosa del Sara	2,16%	San Miguel de Velasco	175
Pailón	2.712.631	4%	Fernández Alonso	1,79%	San Javier	150
Charagua	2.522.711	4%	Santa Cruz de la Sierra	1,76%	San Ignacio de Velasco	145
Santa Rosa del Sara	2.415.050	3%	Cotoca	1,74%	Concepción	142
Yapacaní	2.408.021	3%	Pailón	1,71%	Territorio Indígena Multietnico	129
San Miguel de Velasco	2.248.361	3%	Portachuelo	1,66%	Bella Flor	125
El Carmen Rivero Tórrez	2.238.589	3%	Colpa Bélgica	1,51%	Santa Rosa del Sara	119
San Julián	1.750.913	3%	Mineros	1,45%	San Andrés	107
San Rafael	1.604.501	2%	Tiquipaya	1,27%	Bolpebra	106
San Andrés	1.477.371	2%	Warnes	1,24%	Loreto	103
Santa Ana de Yacuma	1.391.468	2%	El Puente	1,20%	San José de Chiquitos	103
Guayaramerín	1.020.706	1%	San Pedro	1,10%	Ascención de Guarayos	94
San Borja	985.062	1%	San Ramón	0,93%	Baures	93
San Javier	902.678	1%	San Juan	0,91%	Santa Ana de Yacuma	80
Villa Tunari	896.614	1%	San Carlos	0,86%	Ixiamas	75
Puerto Suarez	845.820	1%	Montero	0,85%	Charagua	72
Ixiamas	810.880	1%	Porongo	0,75%	Exaltación	67
Riberalta	807.209	1%	Yacuibá	0,64%	San Joaquín	65
Cuatro Cañadas	803.958	1%	San Javier	0,59%	Reyes	63
Reyes	740.348	1%	La Guardia	0,59%	Pailón	60
San Ignacio	724.790	1%	General Saavedra	0,58%	San Julián	56

Source: Authors' elaboration.

Map 8. Municipalities with higher emissions from deforestation in Bolivia.



Source: Authors' elaboration.

2.5. Sensitivity analysis

The use of average biomass as the basis for biomass loss by forest type represents a key difference from other studies. Default IPCC biomass values for forest types are closer to the 75th percentile of biomass estimates derived from the biomass map used in this study (Santoro & Cartus, 2024). To calculate average biomass, we combined data from intact forests up to 2010, using MapBiomass data to track intact forests from 1985 to 2010. Forests classified as intact until 2010 could be assumed to be at least 25 years old but mature forest could be older but not identifiable from the available data. Given that older forests likely fall within the 75th percentile of biomass content, we conducted an alternative bookkeeping model simulation using this higher biomass estimate.

Using the 75th percentile biomass as the default assumption led to a 23% increase in net emissions from land-cover changes, while the overall composition of net emissions across forest types remained similar. However, it is also possible that deforestation and fires would favour pixels with relatively open forest, which are cheaper to clear and which burn more easily. Thus, we redo the calculations using the 25th percentile biomass value for each forest

type. This assumption implies that less dense forests are deforested first and that newly regrown forests contain lower biomass density than the average. Applying this lower biomass assumption results in a 27% reduction in net emissions. Other modifications were applied to test the sensitivity of the results. Increasing soil carbon loss to 35% over 20 years — assuming more intensive agricultural use — led to a marginal increase of just 2% in net emissions.

3. Discussion

This study aimed to quantify and present CO₂ emissions from deforestation and degradation by forest fires in Bolivia. Separate analyses were conducted, and results are presented independently. To estimate degradation emissions, we subtracted deforestation-related emissions from fire-affected areas. However, integrating both into a single final estimate presents challenges that require careful consideration.

Table 14 shows net emissions from deforestation alongside fire-related emissions, where emissions from deforested areas have been excluded from total fire emissions. Additionally, a “total emissions” category has been included by summing both components. However, these results should be interpreted with caution, particularly regarding fire emissions. This complexity may explain why the Global Carbon Budget (GCB) reports land-use change emissions and fire emissions separately, as they are not directly comparable.

It is important to consider that:

- Degradation emissions account for biomass loss across all vegetation types, not just forests. Most burned areas consist of non-forest vegetation with very low carbon content, except in the Amazónico, Chiquitano, and Chaco Forest zones, where biomass averages ~20 tC/ha.
- Due to explanations exposed in the fire emissions from degradation results, it is very difficult to try to make an assumption of the biomass relationship between forest and non-forest vegetation to disaggregate fire emissions from forest degradation and non-forest vegetation degradation. It would require a more in-depth analysis focused specifically on that topic which goes beyond the scope and duration of the present study.
- Fire emissions reflect gross annual emissions from burning but do not account for subsequent regeneration. Unlike regrowth after deforestation — which can be tracked using land-use change and transition maps — post-fire regeneration is harder to quantify. Estimating net emissions from fires would require annual biomass maps and potentially additional data, such as fire intensity, fire duration, and other variables relevant to both the extent of degradation and post-fire regrowth rates. As shown by exploratory results presented in the fire analysis section, the annual biomass data we have is not enough to support definitive conclusions.


With these considerations in mind, the following table provides an approximate estimate of total annual emissions from both fires and deforestation.

Table 14. Annual fluxes (tCO₂/year) from deforestation and degradation by fire, 2010 – 2022.

Year	Deforestation Net emissions	Estimated emissions by degradation	Total emissions
2010	54.701.914	321.194.068	375.895.982
2011	97.409.559	90.947.327	188.356.887
2012	71.119.561	60.287.903	131.407.463
2013	42.602.309	36.514.693	79.117.002
2014	60.113.008	30.208.427	90.321.435
2015	43.029.928	61.160.012	104.189.940
2016	62.402.727	119.454.330	181.857.057
2017	64.218.445	79.566.359	143.784.803
2018	63.391.560	71.405.972	134.797.532
2019	81.536.224	245.912.314	327.448.539
2020	64.260.985	168.619.422	232.880.408
2021	98.628.832	181.667.620	280.296.452
2022	72.265.981	172.404.627	244.670.609

Source: Authors' elaboration.

Map 9. Average emissions from degradation by fire, 2010–2022.



Source: Authors' elaboration.

Although there is significant year-to-year variation, with 2010 standing out as an extreme year for forest fires, the past four years (2019–2022) have consistently recorded exceptionally high emissions from both deforestation and fire-induced forest degradation, as forest constitute a higher proportion of land cover within burned areas compared to the previous years. Map 9 shows the distribution of the estimated emissions by degradation.

While natural climate variability and climate change, including increasing temperature, shifts in rainfall patterns and wind dynamics, will continue to drive fluctuations in deforestation and fire activity, Bolivia's forests are now more fragmented and frequently burned than ever before. As a result, they have become increasingly vulnerable to external threats, making them more susceptible to future degradation and carbon losses than in the past.

4. Conclusions

Deforestation rates in Bolivia remain high and continue to rise, leading to a corresponding increase in carbon emissions. The majority of emissions due to forest loss are directly attributable to anthropogenic activities, and even the smaller remaining portion cannot be discarded as influenced by human-driven factors.

The highest emissions are concentrated in the department of Santa Cruz, though recent trends show a slight geographical expansion compared to previous decades. Among the top 25 municipalities contributing the most to national emissions, most are located in Santa Cruz and Beni, with one in La Paz and another in Cochabamba. This spatial variability can be explained by two main factors. First, most of the municipalities with the highest emissions have experienced rapid agricultural expansion in recent years. Second, many of these municipalities are located within the Amazon and Chiquitano forests, which have a higher average biomass content per hectare compared to other forest types.

While deforestation emissions are highly attributable to land-use change, fires affect much larger areas, and in most cases, no clear land-use change is observed following the fires. Between 2010 and 2023, Bolivia lost over 5 million hectares of forest, leading to average annual net emissions of around 70 million tons of CO₂ from deforestation. In addition, fire-related gross emissions from forest and non-forest vegetation degradation averaged 126 million tons of CO₂ per year. On a per capita basis, this translates to an annual average of approximately 6,6 tCO₂/person/year from deforestation and 12,3 tCO₂/person/year from fire-related degradation — emission levels that are high compared to those of other sectors and countries. For instance, the global average per capita emissions from land use change are around 1,4 tCO₂/year and in Brazil, the global leader in land-use change emissions (Global Carbon Atlas, 2023), the net emissions from land use change taking into account its large extension is 0,97 tCO₂/ha/year, while Bolivia's reach 0,63 tCO₂/ha/year, a relatively close and concerning figure given the vast difference in country size. Moreover, Brazil's net per capita emissions from deforestation average 4 tCO₂/person/year, whereas Bolivia's 6,6 tCO₂/person/year place it above that level. Only a few countries, such as Belize (11,3 tCO₂/person/year), exhibit higher per capita emissions. Another point of comparison is the reported emissions from Maersk, one of the largest shipping companies in the world, which reported emitting 83,5 million tons of CO₂ in 2024 (Maersk, 2025). In contrast, Bolivia's annual average emissions amount to about 70 million tons of CO₂ from deforestation, plus an additional 126 million tons from fires.

The different forest types present in Bolivia lead to variations in carbon emissions and removals. Forests such as the Amazonian, Yungas, and Chiquitano show high potential for both carbon absorption and emissions, while the Chaco forest currently stands out as a major emitter due to its large spatial extent of affected forest. This study provides guidance for prioritizing areas of focus. For example, the municipalities with the highest contributions to national emissions – such as San Ignacio de Velasco, El Puente, and San José de Chiquitos, along with the top 25 municipalities ranked under each criterion – are mostly located within the Amazon and Chiquitano forests. The latter, in particular, is a high-biomass-density forest found almost entirely in the department of Santa Cruz, and it represents the largest contributor to emissions in the country, both from deforestation and fire-induced degradation. In addition, forest regrowth is less common in this region compared to the Amazon. Therefore, the municipalities with the highest emissions and the two most affected forest types, particularly the Chiquitano forest, should be prioritized to define an effective path toward reducing deforestation and fire rates. However, for smaller areas of interest, more detailed studies are recommended – particularly those incorporating field-based biomass measurements – for forest types with highly variable biomass content.

The estimates presented in this research represent one of the first efforts to disaggregate emissions from deforestation and forest degradation in Bolivia. However, a degree of uncertainty remains. Fire-related data, in particular, offers significant opportunities for further exploration and could be examined in greater depth in future analyses. For instance, further analysis of fire-related data – considering forest types, land cover categories, fire recurrence, and more detailed breakdowns – could enhance our understanding of fire emission dynamics. This includes explaining patterns such as the relatively low emissions from fire-related deforestation – partly due to the fact that about one-third of this deforestation occurs in the Chaco region, which has lower biomass density. Improving the understanding of fire-related emissions would require an improvement of the spatial resolution of available fire emissions and burned area, that are typically available at a resolution of 500x500 meters only compared to 30x30 meters for land cover maps. Despite these limitations this study presents a highly disaggregated spatial analysis with greater resolution and precision than some global estimates. This approach could be replicated, particularly in other South American countries where the MapBiomass project has generated similar high-resolution, temporally detailed land cover data.

The findings confirm that forest loss plays a major role in contributing to climate change, especially in sparsely populated and low-production countries such as Bolivia. This highlights the need to move beyond and to conduct further research aimed at understanding the causes and socioeconomic dynamics behind forest loss, in order to

better address it and develop realistic solutions. More generally, there is a growing need to better understand fire emissions, which also appear to generate significant contributions to global greenhouse gas emissions. This issue is further exacerbated by prolonged dry seasons and other climate-related factors, suggesting a positive feedbacks with higher fire emissions as global warming intensifies.

Bibliography

ANDERSEN, L., ARGANDOÑA, F., CALDERÓN, D., CHOQUE, S., MUÑOZ, Á., OLMOS, C., & MIRANDA, S. (2025).

Valoración económica de los servicios ecosistémicos provistos por las áreas naturales, Áreas Protegidas y los Territorios Indígenas en Bolivia. *La Paz: SDSN Bolivia*. Retrieved from <https://sdsnbolivia.org/documento-de-trabajo-n-1-2025/>

ANDERSEN, L., DOYLE, A. S., DEL GRANADO, S., LEDEZMA, J., MEDINACELI, A., & WEINHOLD, D. (2016). Net Carbon Emissions from Deforestation in Bolivia during 1990–2000 and 2000–2010: Results from a Carbon Bookkeeping Model. *PLOS ONE*, 11(3). <https://doi.org/10.1371/journal.pone.0151241>

APMT. (2020). Tercera Comunicación Nacional del Estado Plurinacional de Bolivia. *Ministerio de Medio Ambiente y Agua*.

CAMBRIDGE UNIVERSITY PRESS. (N.D.). Deforestation. In *Cambridge Dictionary*. Retrieved from <https://dictionary.cambridge.org/dictionary/english/deforestation>

FAO (2018). Global Soil Organic Carbon map (GSOCmap). Rome: FAO.

FAO (2023). “Terms and definitions FRA 2025” Forest Resources Assessment Working Paper 194. Rome: Food and

Agriculture Organization of the United Nations. Retrieved from <https://www.fao.org/3/cc4691en/cc4691en.pdf>

FAO & ITPS. (2020). Global Soil Organic Carbon Map V1.5: Technical Report. Rome: Food and Agriculture Organization of the United Nations.

FRIEDLINGSTEIN, P., O’SULLIVAN, M., JONES, M., ANDREW, R., & BAKKER, D. ET AL. (2023). Global Carbon Budget 2023. *Earth System Science Data*. <https://doi.org/10.5194/essd-15-5301-2023>

GCB (2025). The Global Carbon Budget: FAQs. Retrieved from <https://globalcarbonbudget.org/faqs/>

GLOBAL FOREST WATCH. (N.D.). Tree cover loss in Brazil. Retrieved November 7, 2025, from <https://www.globalforestwatch.org>

GIGLIO, L., JUSTICE, C., BOSCHETTI, L., & ROY, D. (2019). MODIS/Terra+Aqua Burned Area Monthly L3 Global 500m SIN Grid V061. <https://doi.org/10.5067/MODIS/MC D64A1.061>

GLOBAL CARBON ATLAS (2023). Retrieved from <https://globalcarbonatlas.org/>

HANSIS, E., DAVIS, S., & PONG, J. (2015). Relevance of methodological choices for accounting of land use change

carbon fluxes. *Global Biogeochemical Cycles*, 29, 1230–1246. <https://doi.org/doi:10.1002/2014GB 004997>.

HOUGHTON, R., HOBBIE, J., MELILLO, J., MOORE, B., PETERSON, B., SHAVER, G., & WOODWELL, G. (1983). Changes in the Carbon Content of Terrestrial Biota and Soils between 1860 and 1980: A Net Release of CO₂ to the Atmosphere. *Ecological Monographs*, 53(3), 235–262. <https://doi.org/10.2307/1942531>

HOUGHTON, R., & HACKLER, J. (2001). Emissions of carbon from forestry and land-use change in tropical Asia. *Global Change Biology*, 5(4), 481–492. <https://doi.org/10.1046/j.1365-2486.1999.00244.x>

HOUGHTON, R., & CASTANHO, A. (2023). Annual emissions of carbon from land use, land-use change, and forestry from 1850 to 2020. *Earth System Science Data*, 15(5), 2025–2054. <https://doi.org/10.5194/essd-15-2025-2023>

IBISCH, P., & MERIDA, G. (2003). Biodiversidad: La Riqueza de Bolivia. Estado de Conocimiento y Conservación. Santa Cruz de la Sierra: FAN.

IPCC (2006). 2006 IPCC Guidelines for National Greenhouse Gas Inventories. S. Eggleston, L. Buendia, K. Miwa, T.

Ngara, & K. Tanabe, editors.
IPCC/IGES. https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/0_Overview/V0_0_Cover.pdf

IPCC (2019). 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Buendia, E. C., Tanabe, K., Kranjc, A., Jamsranjav, B., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P. & Frederici, S. editors.
<https://www.ipcc-nggip.iges.or.jp/public/2019rf/vol4.html>

IPCC (2024). Report of the IPCC Expert Meeting on Reconciling Anthropogenic Land Use Emissions. Eds: Enoki T., Hayat, M., Grassi G., Sanz M., Rojas Y., Federici S., Seneviratne S., Rupakheti M., Howden M., Sukumar R., Fuglestvedt J., Itsoua Madzous G., Krug T., Romanowskaya A. Pub. IGES, Japan. Retrieved from https://www.ipcc-nggip.iges.or.jp/public/mtdocs/pdfiles/2407_EM_Land_Report.pdf

LANLY, J. (2003). Los factores de la deforestación y la degradación de los bosques. Retrieved from Organización de las Naciones Unidas para la Agricultura y la Alimentación (FAO): <https://www.fao.org/4/xii/ms12a-s.htm>

MAERSK. (2025). Annual report 2024.
<https://www.maersk.com/sustainability/highlights-2024>

MAILLARD, O. (2023). Post-Fire Natural Regeneration Trends in

Bolivia: 2001–2021. *Fire*, 6(1) 18.
<https://doi.org/10.3390/fire6010018>

MAPBIOMAS (2024). ATBD – STEP BY STEP. Retrieved from <https://bolivia.mapbiomas.org/en/atbd-entienda-cada-etapa/>

MAPBIOMAS Bolivia (2024). Colección 2 de la Serie anual de Mapas de Cobertura y Uso del Suelo de Bolivia. Retrieved from <https://bolivia.mapbiomas.org/>

MMAYa (2022). Memoria Técnica: Mapa de bosque 2022. Memoria Técnica, Ministerio de Medio Ambiente y Agua, Viceministerio de Medio Ambiente, Biodiversidad, Cambios Climáticos y de Gestión y Desarrollo Forestal. Retrieved from <https://datos.siarh.gob.bo/biblioteca/686>

POULTER, B., CANADELL, J., HAYES, D., & THOMPSON, R. (2022). Balancing Greenhouse Gas Budgets: Accounting for Natural and Anthropogenic Flows of CO₂ and other Trace Gases. In B. Poulter, A. Bastos, J. Canadell, P. Ciais, D. Huntzinger, R. Houghton, & W. Kurz, Bottom-up approaches for estimating terrestrial GHG budgets: Bookkeeping, process-based modeling, and data-driven methods (pp. 59–85). Amsterdam: Elsevier.

QIN, Z., ZHU, Y., CANADELL, J., CHEN, M., LI, T., MISHRA, U., & YUAN, W. (2024). Global spatially explicit carbon emissions from land use change over the past six decades (1961–2020). *One Earth*, 7, 835–847.

<https://doi.org/10.1016/j.oneear.2024.04.002>

RAISG (2024). Manual General MapBiomas. Retrieved from MapBiomas Bolivia: https://bolivia.mapbiomas.org/wp-content/uploads/sites/15/2024/07/ATBD_General_MapBiomas_Bolivia_-_Coleccion_2.pdf

SANTORO, M., & CARTUS, O. (2024). ESA Biomass Climate Change Initiative (Biomass_cci): Global datasets of forest above-ground biomass for the years 2010, 2015, 2016, 2017, 2018, 2019, 2020 and 2021, v5.01. NERC EDS Centre for Environmental Data Analysis.
<https://doi.org/doi:10.5285/bf535053562141c6bb7ad831f5998d77>

SIMS, M., GIBBS, D., & HARRIS, N. (2024). Greenhouse Gas Fluxes from Forests. Retrieved from World Resources Institute Global Forest Review: <https://gfr.wri.org/biodiversity-ecological-services-indicators/greenhouse-gas-fluxes-forests>

VAN WEES, D., VAN DER WERF, G. R., T. RANDERSON, J., ROGERS, B. M., CHEN, Y., VERAVERBEKE, S., ... MORTON, D. C. (2022). Global biomass burning fuel consumption and emissions at 500 m spatial resolution based on the Global Fire Emissions Database (GFED). *Geoscientific Model Development*, 15, 8411/8437.
<https://doi.org/10.5194/gmd-15-8411-2022>

VIGLIONE, G. (2023). Wildfires. Retrieved from Q&A: How

scientists tackle the challenges
of estimating wildfire CO2
emissions:
<https://www.carbonbrief.org/qa-how-scientists-tackle-the-challenges-of-estimating-wildfire-co2-emission>

What is AFD?

Éditions Agence française de développement publishes analysis and research on sustainable development issues. Conducted with numerous partners in the Global North and South, these publications contribute to a better understanding of the challenges faced by our planet and to the implementation of concerted actions within the framework of the Sustainable Development Goals.

With a catalogue of more than 1,000 titles and an average of 80 new publications published every year, Éditions Agence française de développement promotes the dissemination of knowledge and expertise, both in AFD's own publications and through key partnerships. Discover all our publications in open access at editions.afd.fr.

Towards a world in common.

Publication Director Rémy Rioux

Editor-in-Chief Thomas Melonio

Legal deposit 3rd quarter 2025

ISSN 2492 – 2846

Rights and permissions

Creative Commons license

Attribution – No commercialization – No modification

<https://creativecommons.org/licenses/by-nc-nd/4.0/>



Graphic design MeMo, Juliegilles, D. Cazeils

Layout PUB

Printed by the AFD reprography service

To browse our publications:

<https://www.afd.fr/en/ressources-accueil>